

Warmup Slip

Directions: Turn the following into balanced equations by filling in the blanks with the correct coefficients, formulas of ions or solids, and names.

Cation	Anion	Formula	Name
A. <u> </u> Cu^+	+ <u> </u> SO_4^{2-}	\rightarrow _____	_____
B. <u> </u> Pb^{2+}	+ <u> </u> Cl^-	\rightarrow _____	_____
C. _____	+ _____	\rightarrow AlCl_3	_____
D. _____	+ _____	\rightarrow $\text{Mg}(\text{NO}_3)_2$	_____
E. _____	+ _____	\rightarrow _____	iron (III) sulfide
F. _____	+ _____	\rightarrow _____	copper (II) nitrate
G. <u> </u> Cu^{2+}	+ <u> </u> OH^-	\rightarrow _____	_____

Warmup Slip

Directions: Turn the following into balanced equations by filling in the blanks with the correct coefficients, formulas of ions or solids, and names.

Cation	Anion	Formula	Name
A. <u> 2 </u> Cu^+	+ <u> 1 </u> SO_4^{2-}	\rightarrow Cu_2SO_4	<u>copper (I) sulfate</u>
B. <u> 1 </u> Pb^{2+}	+ <u> 2 </u> Cl^-	\rightarrow PbCl_2	<u>Lead (II) chloride</u>
C. <u> Al³⁺ </u>	+ <u> Cl⁻ </u>	\rightarrow AlCl_3	<u>ALUMINUM CHLORIDE</u>
D. <u> Mg²⁺ </u>	+ <u> NO₃⁻¹ </u>	\rightarrow $\text{Mg}(\text{NO}_3)_2$	<u>Magnesium Nitrate</u>
E. <u> Fe⁺³ </u>	+ <u> S²⁻ </u>	\rightarrow Fe_2S_3	iron (III) sulfide
F. <u> Cu²⁺ </u>	+ <u> NO₃⁻¹ </u>	\rightarrow $\text{Cu}(\text{NO}_3)_2$	copper (II) nitrate
G. <u> 1 </u> Cu^{2+}	+ <u> 2 </u> OH^-	\rightarrow $\text{Cu}(\text{OH})_2$	<u>copper hydroxide</u>

Tests back at 11:45

Friday - quiz

Purpose:

If given a solid substance, how many pieces will its aqueous solution have?

WARMUP finish the warmup slip

#1 AQUEOUS SOLUTIONS

Have water as the solvent

#2 SOLUTIONS

dissolved particles are atomic scale; too small to settle out.

#3 DECIDE WHETHER IONIC OR MOLECULAR

SOLUTIONS ARE USUALLY
IONIC IF THEY HAVE

1) Metal + NonMetal

OR

2) COMMON IONS
FROM BACK OF TABLE

examples

CaCl_2 is ionic, makes Ca , Cl , Cl

NH_4Cl is ionic, makes NH_4 , Cl

$\text{C}_6\text{H}_{12}\text{O}_6$ is molecular, makes $\text{C}_6\text{H}_{12}\text{O}_6$


#4 How many pieces?

A) CHOP IN HALF (+) and (-)

B) EACH COMMON ION MAKES
ONE PIECE

C) IF NOT A COMMON ION,
EACH ATOM MAKES ONE PIECE


#5 How to draw dissolved ions in aqueous solutions.



If someone says
"Draw $\text{CaCl}_2(s)$ after
it gets dissolved
in water to form
an aqueous solution"
you should draw a picture
like the one above.



1. What does STP stand for? Standard temperature and pressure
 2. At STP what is the pressure? 1.00 atmospheres
 3. At STP what is the temperature? 0° Celsius
5. Suppose 8.61 g of zinc was allowed to react with 8.61 liters of HCl gas to produce zinc chloride and hydrogen gas.



b. Which reactant is limiting? (Show both calculations.)

$$8.61 \text{ liters HCl} \times \left(\frac{1 \text{ mol HCl}}{22.4 \text{ L HCl}} \right) \times \left(\frac{1 \text{ mol H}_2}{2 \text{ mol HCl}} \right) = 0.192 \text{ mol H}_2$$

$$8.61 \text{ grams Zn} \times \left(\frac{1 \text{ mol Zn}}{65.38 \text{ g Zn}} \right) \times \left(\frac{1 \text{ mol H}_2}{1 \text{ mol Zn}} \right) = 0.132 \text{ mol H}_2$$


* Answer: Zinc is limiting

- c. Using the limiting reactant, solve for how many liters of hydrogen gas will form at Standard Temperature and Pressure.

We just found that the limiting reactant is Zinc. It formed 0.132 moles of H₂ above.

$$0.132 \text{ mol H}_2 \times \left(\frac{22.4 \text{ L H}_2}{1 \text{ mol H}_2} \right) = 2.96 \text{ liters}$$

6. Determine the volume in liters of carbon dioxide that should be produced in the reaction between 100. g of carbon and 100. liters of O₂. (Use the same three steps as in the previous problem above).



$$100 \text{ g C} \left(\frac{1 \text{ moles}}{12.01 \text{ grams}} \right) \left(\frac{1 \text{ moles CO}_2}{1 \text{ moles C}} \right) = 8.32 \text{ moles CO}_2$$

$$100 \text{ L O}_2 \left(\frac{1 \text{ moles}}{22.4 \text{ liters}} \right) \left(\frac{1 \text{ moles CO}_2}{1 \text{ O}_2} \right) = 4.46 \text{ moles CO}_2$$

then convert the CO₂ from the true result (from the limiting reactant)

$$4.46 \text{ moles CO}_2 \times \left(\frac{22.4 \text{ L CO}_2}{1 \text{ mole CO}_2} \right) = 99.9 \text{ liters}$$


Answer: 99.9 liters CO₂

$$4.46 \text{ mol CO}_2 \times \left(\frac{22.4 \text{ liters}}{1 \text{ moles CO}_2} \right) = 99.9 \text{ liters}$$

KEY

- 7 Suppose 2.00 L of nitrogen gas and 5.00 L of hydrogen gas are mixed and reacted to form ammonia (NH₃). Calculate the volume in liters of ammonia produced when this reaction runs to completion.

Step 1: balanced equation:



Step 2: $2.00 \text{ L N}_2 \times \left(\frac{1 \text{ mol N}_2}{22.4 \text{ L N}_2} \right) \times \left(\frac{2 \text{ mol NH}_3}{1 \text{ mol N}_2} \right) \left(\frac{22.4 \text{ L NH}_3}{1 \text{ mol NH}_3} \right) = 4.00 \text{ L NH}_3$


there are multiple ways to solve. Since the answer is supposed to be in liters of NH₃ gas, we can save time by putting that in the goal step 2.

$$5.00 \text{ L H}_2 \times \left(\frac{1 \text{ mol H}_2}{22.4 \text{ L H}_2} \right) \times \left(\frac{2 \text{ mol NH}_3}{3 \text{ mol H}_2} \right) \left(\frac{22.4 \text{ L NH}_3}{1 \text{ mol NH}_3} \right) = 3.33 \text{ L NH}_3$$

Step 3: We sort of already combined step 3 into step 2. That's why we went into liters instead of stopping at moles.

Answer
3.33 liters
of NH₃ gas

- 8 Uranium (III) sulfide can react with fluorine to form uranium hexafluoride gas and S₈. Write a balanced reaction and then find how many liters at STP of UF₆ will form if you have 333.3 grams uranium (III) sulfide and 0.9 liters fluorine.



Step 2+3: $333.3 \text{ grams U}_2\text{S}_3 \times \left(\frac{1 \text{ mole U}_2\text{S}_3}{572.24 \text{ grams U}_2\text{S}_3} \right) \times \left(\frac{16 \text{ moles UF}_6}{8 \text{ moles U}_2\text{S}_3} \right) \left(\frac{22.4 \text{ liters UF}_6}{1 \text{ mole UF}_6} \right) = 26.1 \text{ L UF}_6$

$$0.9 \text{ liters F}_2 \times \left(\frac{1 \text{ mole F}_2}{22.4 \text{ liters F}_2} \right) \times \left(\frac{16 \text{ moles UF}_6}{48 \text{ moles F}_2} \right) \left(\frac{22.4 \text{ liters UF}_6}{1 \text{ mole UF}_6} \right) = 0.3 \text{ L UF}_6$$

↑
use
one mole = 22.4 L

↑
use the coefficients from the balanced equation

↑
use
one = 22.4 L gas at STP

Answer: F₂ is limiting. 0.3 liters of UF₆ will form.